

A**Alexander M.P.**

Clinical determination of mental competence. A theory and a retrospective study.
Arch. Neurol. 1988, 45(1): 23-26.

American Psychiatric Association.

Diagnostic Criteria from DSM III - R. Washington, 1987.

Anthony J.C., Le Resche L., Niaz U., Von Korff M.R., Folstein M.F.

Limits of the "Mini Mental State" as a screening test for dementia and delirium among hospital patients.
Psychol. Med. 1982, 12(2): 397-340.

Ashworth B.

Preliminary trial of carisprodol in multiple sclerosis.
Practitioner 1964, 192 : 540-542.

B**Bellamy N., Buchanan W.W., Goldsmith C.H., Campbell J., Stitt L.W.**

Validation study of WOMAC: a health instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip and knee.
J. Rheumatol. 1988, 15(12): 1833-1840.

Benaïm C., Pérennou D.A., Villy J., Rousseaux M., Pélissier J.Y.

Validation of a standardized assessment of postural control in stroke patients: the Postural Assessment Scale for Stroke Patients (PASS).
Stroke 1999, 30(9): 1862-1868.

Benaïm C., Cailly B., Pérennou D., Pélissier J.

Validation of the Aphasic Depression Rating Scale.
Stroke 2004, 35(7): 1692-1696.

Benaïm C., Froger F., Gazottes C., Gueben D., Porte M., Desnuelle C., Pélissier J.

Use of the Faces Pain Scale by left and right hemispheric stroke patients.
Pain 2007, 128(4): 52-58.

Bérard C., Payan C., Fermanian J., Girardot F., et le groupe d'étude MFM.

La mesure de fonction motrice, outil d'évaluation clinique des maladies neuromusculaires. Étude de validation.
Rev. Neurol. (Paris) 2006, 162(4) : 485-493.

Bergego C., Azouvi P., Samuel C., Marchal F., Louis-Dreyfus A. et al.

Validation d'une échelle d'évaluation fonctionnelle de l'héminégligence dans la vie quotidienne : l'échelle CB.
Ann. Readapt. Med. Phys. 1995, 38 : 183-189.

Bergner M., Bobbitt R.A., Pollard W.E., Martin D.P., Gilson B.S.

The Sickness Impact Profile: validation of a health status measure.
Med. Care 1976, 14 (1): 57-67.

Béthoux F., Calmels P.

Guide des outils de mesure et d'évaluation en médecine physique et de réadaptation.
Éditions Frison-Roche, 2003.

Bieri D., Reeve R.A., Champion G.D., Addicoat L., Ziegler J.B.

The Faces Pain Scale for the self-assessment of the severity of pain experienced by children: development, initial validation, and preliminary investigation for ratio scale properties.
Pain 1990, 41(2): 139-150.

Bocquet H., Pous J., Charlet J.P., Grand A.

Mesure de la charge des soignants chez les personnes âgées dépendantes par l'inventaire de Zarit.
Rev. Épidemiol. Santé Publique 1996, 44(1) : 57-65.

Bohannon R.W., Smith M.B.

Interrater reliability of a modified Ashworth scale of muscle spasticity.
Phys. Ther. 1987, 67(2): 206-207.

Borg A.

Perceived exertion as an indicator of somatic stress.
Scand. J. Rehabil. 1970, 2(2): 92-98.

Born J.D., Hans P., Dexters G., Kalangu K., Lenelle J., Milbouw G., Stevenaert A.

Évaluation pratique du dysfonctionnement encéphalique chez le traumatisé crânien.
Acta Neurochir. (Wien) 1988, 91(1-2): 1-11.

Born J.D.

The Glasgow-Liège Scale. Prognostic value and evolution of motor response and brain stem reflexes after severe head injury.
Muscle Nerve 1983, 6(2): 91-103.

Boureau F., Luu M., Doubrere J.F., Gay C.

Élaboration d'un questionnaire d'auto-évaluation de la douleur par liste de qualitatifs. Comparaison avec le Mac Gill pain questionnaire de Melzack.
Thérapie 1984, 39(2) : 119-129.

Boureau F., Luu M.

Les méthodes d'évaluation de la douleur clinique.
Doul. Analg. 1988, 1 : 65-73.

Boussagol B., Pélissier J., Hérisson C., Simon L.

Évaluation des pathologies de la coiffe des rotateurs : analyse comparative de quatre scores.
Ann. Réadapt. Med. Phys. 1996, 39 : 79-88.

Bowen J., Gibbons L., Gianas A., Kraft G.H.

TSelf-administered Expanded Disability Status Scale with functional system scores correlates well with a physician-administered test.
Mult. Scler. 2001, 7(3): 201-206.

Braden B., Bergstrom N.

A conceptual schema for the study of the etiology of pressure sores.
Rehabil. Nurs. 1987, 12(1): 8-12.

Brooke M.H., Fenichel G.M., Griggs R.C., Mendell J.R., Moxley R. et al.

Clinical investigation in Duchenne dystrophy: 2. Determination of the "power" of therapeutic trials based on natural history.
Muscle Nerve 1983, 6(2): 91-103.

Brooke M.

A clinical's view of neuromuscular disease.
2nd ed. Baltimore : williams & Wilkins ; 1986.

Brott T., Adams H.P. Jr., Olinger C.P., Marler J.R. et al.

Measurements of acute cerebral infarction: a clinical examination scale.
Stroke 1989, 20(7): 864-870.

Brun V., Dhoms G. et al.

L'équilibre posturale de l'hémiplégique : proposition d'indices d'évaluation.
In : Simon L., Pélissier J., Hérisson C. Ed : Actualités en rééducation fonctionnelle et réadaptation..
Paris, Masson : 1991, pp : 412-417.

Brun V., Mousbeh Z., Jouet-Pastre B. et al.

Evaluation clinique de la marche de l'hémiplégique vasculaire : proposition d'une modification de la Functional Ambulation Classification (FAC).
Ann. Réadapt. Med. Phys. 2000, 43 : 14-20.

C

Chaory K., Fayad F., Rannou F., Lefèvre-Colau M.M., Fermanian J. et al.
Validation of the french version of the Fear Avoidance Belief Questionnaire.
Spine 2004, 29(8): 908-913.

Chwalow A.J., Lurie A., Bean K., Parent du Chatelet I., Venot A. et al.
A French version of the Sickness Impact profile (SIP): stages in the cross cultural validation of a generic quality of life scale.
Fundam. Clin. Pharmacol. 1992, 6(7): 319-326.

Collignon P., Giusiano B., Boutin A.M., Combes J.C.
Utilisation d'une échelle d'hétéro-évaluation de la douleur chez le sujet sévèrement polyhandicapé.
Doul. Analg. 1997, 1 : 27-32.

Coste J., Le Parc J.M., berge E., Delacoeuillerie G., Paolaggi J.B.
In : Gauvain-Piquard A., Murat I., Pons G. La douleur chez l'enfant : échelles d'évaluation, traitements médicamenteux.
Spinger Verlag, Paris, 1993 : 11-20.

Collignon P., Giusiano B., Combes J.C.
La douleur chez l'enfant polyhandicapé. In : Ecoffey C, Murat I. La douleur chez l'enfant.
Flammarion Med-Sci, Paris, 1999 pp : 174-178.

Coste J., Le Parc J.M., berge E., Delacoeuillerie G., Paolaggi J.B.
Validation française d'une échelle d'incapacité fonctionnelle pour l'évaluation des lombalgies EIFEL.
Rev. Rhumatol. 1993, 60(5) : 335-341.

Crook T., Bartus R.T., Ferris S.H., White House P., Cohen G.D., Gershon S.
Age-associated memory impairment : proposed diagnostic criteria and measures of clinical change.
Report of a National Institute of Mental Health Work Group.
Dev. Neuropsychol. 1986, 2: 261-276.

D

Dérouesne C., Poitreneau J., Hugonot L. et al.
Le Mini-Mental State Examination (MMSE) : un outil pratique pour l'évaluation de l'état cognitif des patients par le clinicien. Version française consensuelle.
Presse Med. 1999, 28(21) : 1141-1148.

Dick J.P., Guiloff R.J., Stewart A., Blackstock J., Bielawska C. et al.
Mini-mental State examination in neurological patients.
J. Neurol. Neurosurg. Psychiatry 1984, 47(5): 496-499.

Ditunno J.F., Scivoletto G., Patrick M., Biering-Sorensen F., Abel R., Marino R.
Validation of the walking index for spinal cord injury in a US and European clinical population.
Spinal Cord 2008, 46(3): 181-188.

Dubois B., Slachevsky A., Levy R.
Démences de type frontal.
La Lettre du neurologue 1997, (N° Hors série, Avril) : 16-19.

Duruoz M.T., Poiraudeau S., Fermanian J., Menkes C.J., Amor B. et al.
Development and validation of a rhumatoïd hand functional disability scale that assesses functional handicap.
J. Rheumatol. 1996, 23(7): 1167-1172.

Enjalbert M., Pélissier J., Blin D., Lopez F.M., Simon L.
La préhension chez l'hémiplégique. II. Facteurs de pronostic.
Ann. Réadapt. Méd. Phys., 1988, 31 : 155-163.

Eslinger P.J., Damasio A.R., Beton A.L., van Allen M.
Neuropsychologic detection of abnormal mental decline in older persons.
JAMA 1985, 253(5): 670-674.

Eustache F., Viel J.F., Dary M., Thieuw F., Lechevallier B., Signoret J.L.
Comparaison des performances cognitives dans la démence de type Alzheimer selon l'âge d'apparition des premiers symptômes ; in : D. Leys et H. Petit. La maladie d'Alzheimer et ses limites.
Congrès de Psychiatrie et de Neurologie de Langue Française. pp. 299-306. Masson, 1988.

F

Fahn S., Elton R.L.

Unified Parkinson's Disease Rating Scale.

In Fahn S., Marsden C.D., Calne D. and Goldstein M. Ed. Recent development in Parkinson disease. Vol. 2, Florham Park NJ. Macmillan Healthcare Information: 1987, pp:153-163.

Fattal C.

Motor capacities of upper limbs in tetraplegics: a new scale for the assessment of the results of functional surgery on upper limbs.

Spinal Cord 2004, 42(2): 80-90.

Fattal C., Thery J.M., Micallef J.P.

Validation of the motor capacities scale: a specific evaluation of manual abilities in tetraplegics who undergo functional surgery of the upper limbs.

Ann. Readapt. Med. Phys. 2004, 47(8): 537-545.

Fleck M.P., Guelfi J.D., Poirier-Littre M.F., Loo H.

Application of a structure interview guide adapted to 4 depression scales.

Encéphale 1994, 20 (5) : 479-486.

Folstein M.F., Folstein S.E., MacHugh P.R.

"Mini Mental State": a practical method for grading the cognitive state of patients for the clinician.

J. Psychiatr. Res. 1975, 12(3): 189-198.

G

Gaspari M., Roveda G., Scandellari C., Stecchi S.

An expert system for the evaluation of EDSS in multiple sclerosis.

Artif. Intell. Med. 2002, 25(2): 187-210.

Gauthier L., Dehaut F., Joanette Y.

The bells test: a quantitative and qualitative test for visual neglect.

Int. J. Clin. Neuropsychol. 1989, 11: 49-53.

Goldstein L.B., Samsa G.P.

Reliability of the National Institutes of Health Stroke Scale. Extension to Non-Neurologists in the context of a clinical trial.
Stroke 1997, 28(2): 307-310.

Goodglass H., Kaplan E.

The assessment of aphasia and related disorders.

Lea and Fibiger: Philadelphia, 1972.

Goodkin D.E., Cookfair D., Wende K., Bourdette D., Pullicino P. et al.

Inter- and intrarater scoring agreement using grades 1.0 to 3.5 of the Kurtzke Expanded Disability Status Scale (EDSS).
Multiple sclerosis Collaborative Research Group.
Neurology 1992, 42(4): 859-863.

Groupe d'étude MFM.

Escale, Unité de rééducation pédiatrique, centre Hospitalier Lyon Sud, 69495 Pierre-Bénite.

Manuel de l'utilisateur, édition 2004.

Guelfi J.D., Criquillon-Doublet S.

Dépression et syndromes anxiol-dépressifs.

Paris : Ed. Pil et Arrix, 1992.

Guivarc'h S., Bérard C., Calmels P.

Affections neuromusculaires. In : Guide des outils de mesure et d'évaluation en médecine physique et de réadaptation.
F. Béthoux, P. Calmels.

Ed Frison-Roche, 2003. Paris, pp : 269-283.

Guivarc'h S.

"Une échelle internationale de mesure de la fonction motrice pour les maladies neuro-musculaires."

Mémoire de DEA, Université de Bourgogne, 1998.

H

Harwood R.H., Rogers A., Dickinson E., Ebrahim S.

Measuring handicap: the London Handicap Scale, a new outcome measure for chronic disease.
Qual. Health Care 1994, 3(1): 11-16.

Huskisson E.C.

Measurement of pain.
Lancet 1974, 2 (7889): 1127-1131.

I

Itzkovich M., Gelernter I., Biering-Sorensen F., Weeks C., Laramee M.T. et al.

Spinal Cord Independence Measure (SCIM) version III: reliability and validity in a multi-center international study.
Disabil. Rehabil. 2007, 29(24): 1926-1933.

J

Jennet B., Bond M.

Assessment of outcome after severe brain damage: a practical scale.
Lancet 1975, 1(7889): 1127-1131.

K

Kasner S.E., Chalela J.A., Luciano J.M., Cucchiara B.L., Raps E.C. et al.

Reliability and validity of estimating the NIH stroke scale score from medical records.
Stroke 1999, 30(8): 1534-1537.

Kasner S.E., Cucchiara B.L., McGarvey M.L., Luciano J.M., Liebeskind D.S., Chalela J.A.

Modified National Institutes of Health Stroke Scale can be estimated from medical records.
Stroke 2003, 34(2): 568-570.

Klein L.E., Roca R.P., McArthur J., Vogelsang G., Klein G.B. et al.

Diagnosing dementia. Univariate and multivariate analyses of the mental status examination.
J. Am. Geriatr. Soc. 1985, 33(7): 483-488.

Kopec J.A., Esdaile J.M., Abrahamowicz M., Abenaim L., Wood-Dauphinee S. et al.

The Quebec back pain disability scale. Measurement properties.
Spine 1995, 20(3): 341-352.

Kopec J.A., Esdaile J.M., Abrahamowicz M., Abenaim L., Wood-Dauphinee S. et al.

The Quebec Back Pain Disability Scale: conceptualization and development.
J. Clin. Epidemiol. 1996, 49(2): 151-161.

Krupp L.B., LaRocca N.G., Muir-Nash J., Steinberg A.D.

The fatigue severity scale. Application to patients with multiple sclerosis and systemic lupus erythematosus.
Arch. Neurol. 1989, 46(10): 1121-1123.

Kuntzer T., Decosterd I.

Douleurs neuropathiques : contexte, nouveaux outils, nouveaux médicaments.
Rev. Med. Suisse. 2005, 1(43) : 2812-2816.

Kurtzke J.F.

Rating neurological impairment in multiple sclerosis: an expanded disability status scale (EDSS).
Neurology 1983, 33(11): 1444-1452.

L

Lacote M., Chevalier A.M., Miranda A., Bleton J.P.

Évaluation Clinique de la fonction musculaire.
3^e édition ; Maloine, Paris, 1996.

Lawlis G.F., Cuencas R., Selby D., McCoy C.E.

The development of the Dallas Pain Questionnaire. An assessment of the impact of spinal pain on behavior.
Spine 1989, 14(5): 511-516.

Leplège A., Ecosse E., Verdier A., Perneger T.V.

The French SF-36 health Survey: translation, cultural adaptation and preliminary psychometric evaluation.
J Clin. Epidemiol. 1998, 51(11): 1013-1023.

Leplège A.

Le questionnaire MOS SF-36, manuel de l'utilisateur et guide d'interprétation des scores.
Ed : Estem, 2001.

Lequesne M., Samson M., Gerard P., Mery C.

Pain-function indices for the follow-up of osteoarthritis of the hip and knee.
Rev. Rhum. Mal. Osteoartic. 1990, 57(9 pt2) : 32S-36S.

Levin H.S., O'Donnell V.M., Grossman R.G.

www.amedoc.com/goat.htm

Levin H.S., O'Donnell V.M., Grossman R.G.

The Galveston Orientation and Amnesia Test.
A practical scale to assess cognition after head injury.
J. Nerv. Ment. Dis. 1979, 167(11): 675-684.

Levin H.S., High W.M., Goethe K.E., Sisson R.A., Overall J.E. et al.

The neurobehavioural rating scale: assessment of the behavioural sequelae of head injury by the clinician.
J. Neurol. Neurosurg. Psychiatry 1987, 50(2): 183-189.

Lundin-olsson L., Nyberg L., Gustafson Y.

"Stops walking when talking" as a predictor of falls in elderly people.
Lancet 1997, 349(9052): 617.

Lyle R.C.

A performance test for assessment of upper limb function in physical rehabilitation treatment and research.
Int. J. Rehabil. Res. 1981, 4(4): 483-492.

M**McDowell C.L., Moberg E.A., Smith A.G.**

International conference on surgical rehabilitation of the upper limb in tetraplegia.
J. Hand Surg. [Am]. 1979, 4(4): 387-390.

Mc Horney C.A., Ware J.E. jr., Raczeck A.E.

The MOS 36-item short-form health survey (SF-36). II. Psychometric and clinical tests of validity in measuring physical and mental headline constructs.
Med. Care 1993, 31(3): 247-263.

Mahoney F.I., Barthel D.W.

Functional evaluation: the Barthel Index.
Md. State Med. J. 1965, 14: 61-65.

Marino R.J., Shea J.A., Stineman M.G.

The Capabilities of Upper Extremity instrument: reliability and validity of a measure of functional limitation in tetraplegia.
Arch. Phys. Med. Rehabil. 1998, 79(12): 1512-1521.

Marty M., Blotman F., Avouac B., Rozenberg S., Valat J.P.

Validation de la version française de l'échelle de Dallas chez les patients lombalgiques chroniques.
Rev. Rhum. 1998, 65(2) : 139-147.

Mathias S., Nayak U.S.L., Isaacs B.

Balance in elderly patients: the "Get-up and Go test".
Arch. Phys. Med. Rehabil. 1986, 67(6): 387-389.

Mathiowetz V., Weber K., Kashman N., Volland G.

Adult norms for the nine-hole peg test of finger dexterity.
Occup. Ther. J. Res. 1985, 39: 386-391.

Mazaux J.M., Dehail P., Brun V., Pélissier J.

Les principaux tests francophones d'évaluation de l'aphasie.
In: J.M. Mazaux, V. Brun et J. Pélissier, Ed. Aphasic Masson Ed., Paris 2000, pp : 78-88.

Minaire P.

La mesure de l'indépendance fonctionnelle (MIF). Historique, présentation et perspectives.
J. Réadapt. Med. 1991, 11 : 168-174.

Morris R.G., Kopelman M.D.

The memory deficit in alzheimer - type Dementia : a review.
Q. J. Exp. Psychol. Am. 1986, 38(4): 575-602.

O**Odderson I.R.**

The National Institutes of Health Stroke Scale and its importance in acute stroke management.
Phys. Med. Rehabil. Clin. N. Am. 1999, 10(4): 787-800.

P**Penn R.D., Savoy S.M., Corcos D., Latash M., Gottlieb G. et al.**

Intrathecal baclofen for sever spinal spasticity.
N. Engl. J. Med. 1989, 320 (23): 1517-1521.

Perneger T.V., Leplège A., Etter J.F., Rougemont A.

Validation of a French-langage version of the MOS 36-Item Short Form Health Survey (SF-36) in young healthy adults.
J. Clin. Epidemiol. 1995, 48(8): 1051-1060.

R**Rousseaux M., Beis J.M., Pradat-Diehl P., Martin Y., Bartolomeo P. et al.**

Mémoire. Présentation d'une batterie de dépistage de la négligence spatiale
Normes et effets de l'âge, du niveau d'éducation, du sexe, de la main et de la latéralité.
Rev. Neurol. (Paris) 2001, 157(11Pt1) : 1385-1400.

Russell D., Rosenbaum P., Cadman D., Gowland C., Hardy S., Jarvis S.

The Gross Motor Function Measure: a mean to evaluate the effects of physical therapy.
Dev. Med. Child Neurol. 1989, 31(3): 341-352.

S**Scott O.M., Hyde S.A., Goddard C., Dubowitz V.**

Quantitation of muscle function in children: a prospective study in Duchenne muscular dystrophy.
Muscle Nerve 1982, 5(4): 291-301.

Shiel A., Horn S.A., Wilson B.A., Watson M.J., Campbell M.J. et al.

The Wessex Head Injury Matrix (WHIM) main scale: a preliminary report on a scale to assess and monitor patient recovery after severe head injury.
Clin. Rehabil. 2000, 14(4): 408-416.

Signoret J.L., Bonvarlet M., Benoit N., Bolgert F., Eustache F., Leger J.M.

Batterie d'estimation des états démentiels ; description et validation ; in : D. Leys et H. Petit. La maladie d'Alzheimer et ses limites. Congrès de Psychiatrie et de Neurologie de Langue Française. pp : 265-270. Masson, 1988.

Smith R.A., Newcombe R.G., Sibert J.R., Harper P.S.

Assessment of locomotor function in young boys with Duchenne muscular dystrophy.
Muscle Nerve 1991, 14(5): 462-469.

Smith T., Gildeh N., Holmes C.

The Montreal Cognitive Assessment: validity and utility in a memory clinic setting.
Can. J. Psychiatry 2007, 52(5): 329-323.

T**Tariot P.N., Weingartner H.**

A psychologic analysis of cognitive failures. Structure and mechanisms.
Arch. Gen. Psychiatry 1986, 43(12):1183-1188.

Teasdale G., Jennett B.

Assessment of coma impaired consciousness. A practical scale.
Lancet 1974, 2(7872): 81-84.

Tinetti M.E.

Performance-oriented assessment of mobility problems in elderly patients.
J. Am. Geriatr. Soc. 1986, 34(2): 119-126.

V

Vernay D., Gerbeaud L., Biolay S., Coste J., Debourse J. et al.

Qualité de vie et sclérose en plaque : validation de la version francophone d'un auto-questionnaire, le SEP-59.
Rev. Neurol. 2000, 156(3) : 247-263.

Vignos P.J. Jr, Spencer G.E. Jr, Archibald K.C.

Management of progressive muscular dystrophy in childhood.
JAMA 1963, 184: 89-96.

W

Waddell G., Newton M., Henderson I., Somerville D., Main C.J.

Fear Avoidance Belief Questionnaire (FABQ) the role of fearavoidance belief in chronic low back pain and disability.
Pain 1993, 52(2): 157-168.

Wade D.T., Langton-Hewer R., Wood V.A., Skilbeck C.E., Ismail H.M.

The hemiplegic arm after stroke: measurement and recovery.
J. Neurol. Neurosurg. Psychiatry 1983, 46(6): 521-524.

Wade D.T.

Measurment in neurological rehabilitation.
New York : Oxford University Press, 1992.

Ware J.E. jr., Sherbourne C.D.

The MOS 36-item short-form health survey (SF-36).
I. Conceptual framework and item selection.
Med. Care 1992, 30(6): 473-483.

Ware J.E. jr.

SF-36 health survey: manual and interpretation guide.
Nimrod press. Boston, 1997.

Wary J.E., Villard J.F.

Doloplus : une échelle d'évaluation de la douleur de la personne âgée.
Le généraliste et la gérontologie 1995, 14 : 26-33.

Williams J.B.

A structure interview guide for Hamilton Depression Rating Scale.
Arch. Gen. Psychiatry 1988, 45(8): 742-747.

Z

Ziegler M.

UPDRS Résumé (version 1985).
Hôpital Léopold Bellan, Unité James Parkinson (Vs : 12/12/97).